

Chickenpox Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Chickenpox (also called varicella) is a common childhood disease. It is usually mild, but it can be serious, especially in young infants and adults.

- It causes a rash, itching, fever, and tiredness.
- It can lead to severe skin infection, scars, pneumonia, brain damage, or death.
- The chickenpox virus can be spread from person to person through the air, or by contact with fluid from chickenpox blisters.
- A person who has had chickenpox can get a painful rash called shingles years later.
- Before the vaccine, about 11,000 people were hospitalized for chickenpox each year in the United States.
- Before the vaccine, about 100 people died each year as a result of chickenpox in the United States.

Chickenpox vaccine can prevent chickenpox.

Most people who get chickenpox vaccine will not get chickenpox. But if someone who has been vaccinated does get chickenpox, it is usually very mild. They will have fewer blisters, are less likely to have a fever, and will recover faster.

2 Who should get chickenpox vaccine and when?

Routine

Children who have never had chickenpox should get 2 doses of chickenpox vaccine at these ages:

- | | |
|-----------|----------------------------------------------------------------------------------|
| 1st Dose: | 12–15 months of age |
| 2nd Dose: | 4–6 years of age (may be given earlier, if at least 3 months after the 1st dose) |

People 13 years of age and older (who have never had chickenpox or received chickenpox vaccine) should get two doses at least 28 days apart.

Catch-up

Anyone who is not fully vaccinated, and never had chickenpox, should receive one or two doses of chickenpox vaccine. The timing of these doses depends on the person's age. Ask your doctor.

Chickenpox vaccine may be given at the same time as other vaccines.

Note: A “combination” vaccine called **MMRV**, which contains both chickenpox and MMR vaccines, may be given instead of the two individual vaccines to people 12 years of age and younger.

3 Some people should not get chickenpox vaccine or should wait.

- People should not get chickenpox vaccine if they have ever had a life-threatening allergic reaction to a previous dose of chickenpox vaccine or to gelatin or the antibiotic neomycin.
- People who are moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting chickenpox vaccine.
- Pregnant women should wait to get chickenpox vaccine until after they have given birth. Women should not get pregnant for 1 month after getting chickenpox vaccine.
- Some people should check with their doctor about whether they should get chickenpox vaccine, including anyone who:
 - Has HIV/AIDS or another disease that affects the immune system
 - Is being treated with drugs that affect the immune system, such as steroids, for 2 weeks or longer
 - Has any kind of cancer
 - Is getting cancer treatment with radiation or drugs
- People who recently had a transfusion or were given other blood products should ask their doctor when they may get chickenpox vaccine.

Ask your doctor for more information.



4**What are the risks from chickenpox vaccine?**

A vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of chickenpox vaccine causing serious harm, or death, is extremely small.

Getting chickenpox vaccine is much safer than getting chickenpox disease. Most people who get chickenpox vaccine do not have any problems with it. Reactions are usually more likely after the first dose than after the second.

Mild problems

- Soreness or swelling where the shot was given (about 1 out of 5 children and up to 1 out of 3 adolescents and adults)
- Fever (1 person out of 10, or less)
- Mild rash, up to a month after vaccination (1 person out of 25). It is possible for these people to infect other members of their household, but this is extremely rare.

Moderate problems

- Seizure (jerking or staring) caused by fever (very rare).

Severe problems

- Pneumonia (very rare)

Other serious problems, including severe brain reactions and low blood count, have been reported after chickenpox vaccination. These happen so rarely experts cannot tell whether they are caused by the vaccine or not. If they are, it is extremely rare.

Note: The first dose of **MMRV** vaccine has been associated with rash and higher rates of fever than MMR and varicella vaccines given separately. Rash has been reported in about 1 person in 20 and fever in about 1 person in 5.

Seizures caused by a fever are also reported more often after MMRV. These usually occur 5–12 days after the first dose.

5**What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS is only for reporting reactions. They do not give medical advice.

6**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

7**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
Varicella Vaccine

3/13/2008

42 U.S.C. § 300aa-26

Office Use Only

